Cambridge IGCSE[™]

| CANDIDATE NAME | | | | | |
|-------------------|--|--|---------------------|--|--|
| CENTRE NUMBER | | | CANDIDATE NUMBER | | |

0123456789

CHEMISTRY 0620/05

Paper 5 Practical Test

For examination from 2023

SPECIMEN PAPER

1 hour 15 minutes

You must answer on the question paper.

You will need: The materials and apparatus listed in the confidential instructions

INSTRUCTIONS

- Answer all questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for each question or part question is shown in brackets [].
- Notes for use in qualitative analysis are provided in the question paper.

| For Examiner's Use | | |
|--------------------|--|--|
| 1 | | |
| 2 | | |
| 3 | | |
| Total | | |

This document has 10 pages. Any blank pages are indicated.

© UCLES 2020 [Turn over

1 You are going to investigate the reaction between dilute hydrochloric acid and two different aqueous solutions of sodium hydroxide labelled solution **A** and solution **B**.

Read all of the instructions carefully before starting the experiments.

Instructions

You are going to do **two** experiments.

(a) Experiment 1

- Rinse a burette with dilute hydrochloric acid.
- Fill the burette with dilute hydrochloric acid.
- Run some of the dilute hydrochloric acid out of the burette so that the level of dilute hydrochloric acid is on the burette scale.
- Record the initial burette reading in Table 1.1.
- Use a measuring cylinder to pour 25 cm³ of solution **A** into a conical flask.
- Add five drops of thymolphthalein indicator to the conical flask.
- While swirling the conical flask, slowly add the dilute hydrochloric acid from the burette to the flask until the solution just changes colour.
- Record the final burette reading in Table 1.1 and complete the table.

Experiment 2

- Fill the burette with dilute hydrochloric acid.
- Run some of the dilute hydrochloric acid out of the burette so that the level of dilute hydrochloric acid is on the burette scale.
- Record the initial burette reading in Table 1.1.
- Empty the conical flask and rinse it with distilled water.
- Use the measuring cylinder to pour 25 cm³ of solution **B** into the conical flask.
- Add five drops of thymolphthalein indicator to the conical flask.
- While swirling the conical flask, slowly add the dilute hydrochloric acid from the burette to the flask until the solution just changes colour.
- Record the final burette reading in Table 1.1 and complete the table.

Table 1.1

| | Experiment 1 | Experiment 2 |
|--|--------------|--------------|
| final burette reading / cm ³ | | |
| initial burette reading / cm ³ | | |
| volume of dilute hydrochloric acid added / cm ³ | | |

| ı | 41 |
|---|----|
| ı | 41 |

(b) State the colour change observed in Experiment 1.

from to [1]

| (c) | (i) | State which solution of sodium hydroxide, solution A or solution B , is the more concentrated. | | | | |
|-----|------|--|--|--|--|--|
| | | Explain your answer. | | | | |
| | | [1] | | | | |
| | (ii) | Deduce the simplest whole number ratio of concentration of solution ${\bf A}$: concentration of solution ${\bf B}$. | | | | |
| | | [1] | | | | |
| (d) | | te the volume of hydrochloric acid needed if Experiment 1 is repeated using $10\mathrm{cm}^3$ of ition A . | | | | |
| | | [2] | | | | |
| (e) | In E | experiment 2 the conical flask is rinsed with distilled water. | | | | |
| | (i) | Suggest why the conical flask is rinsed with distilled water. | | | | |
| | | [1] | | | | |
| | (ii) | The conical flask is not dried after it is rinsed with distilled water. | | | | |
| | | Suggest why the conical flask is not dried. | | | | |
| | | [1] | | | | |
| (f) | | te the effect, if any, on the volume of dilute hydrochloric acid used in Experiment 1 if the ition of sodium hydroxide is warmed before adding the dilute hydrochloric acid. | | | | |
| | Giv | e a reason for your answer. | | | | |
| | effe | ct on volume | | | | |
| | reas | son[2] | | | | |
| (g) | (i) | Suggest how the reliability of the results from Experiment 1 and Experiment 2 can be confirmed. | | | | |
| | | [1] | | | | |
| | (ii) | Suggest a more accurate method of measuring the volume of the solution of sodium hydroxide. | | | | |
| | | [1] | | | | |

barium hydroxide.

(h) Aqueous sodium hydroxide reacts with aqueous barium chloride to form a white precipitate of

| Use this information to suggest a different method of finding out which of the solutions of sodium hydroxide, solution A or solution B , is more concentrated. |
|--|
| In your answer, state how your results show which solution of sodium hydroxide, solution ${\bf A}$ or solution ${\bf B}$, is more concentrated. |
| |
| |
| |
| [3] |
| [Total: 18] |

BLANK PAGE

| Υοι | u are provided with two solids, solid C and solid D . |
|-------------|--|
| Do | the following tests on solid C and solid D , recording all of your observations at each stage. |
| tes | ts on solid C |
| (a) | Describe the appearance of solid C . |
| | [1] |
| (b) | Place about half of solid C in a hard-glass test-tube. Heat the solid gently then strongly. |
| | Record your observations. |
| | |
| | [2] |
| | d the rest of solid C to about 10cm^3 of distilled water in a boiling tube. Stopper the boiling tubed shake it to dissolve solid C and form solution C . |
| Div | ide solution C into four approximately equal portions in four test-tubes. |
| (c) | Test the pH of the first portion of solution C . |
| | pH =[1] |
| (d) | To the second portion of solution C , add about 1 cm depth of dilute nitric acid followed by a few drops of aqueous silver nitrate. |
| | Record your observations. |
| | [1] |
| , , | |
| (e) | To the third portion of solution C , add about 1 cm depth of dilute nitric acid followed by a few drops of aqueous barium nitrate. |
| | Record your observations. |
| | [1] |
| /£ \ | To the fourth parties of solution C add aqueous ammenia dresswice and then in excess |
| (f) | To the fourth portion of solution C , add aqueous ammonia dropwise and then in excess. |
| | Record your observations. |
| | |
| | [2] |

| (g) | Identify solid C . [2] |
|-----|--|
| tes | ts on solid D |
| (h) | Do a flame test on solid D . |
| | Record your observations. |
| | [1] |
| | If the rest of solid $\bf D$ to about 10 cm ³ of distilled water in a boiling tube. Stopper the boiling tube shake it to dissolve solid $\bf D$ and form solution $\bf D$. |
| Div | de solution D into two approximately equal portions in two test-tubes. |
| (i) | To the first portion of solution D , add aqueous sodium hydroxide dropwise and then in excess. |
| | Record your observations. |
| | |
| | [2] |
| (j) | To the second portion of solution D , add about 1 cm depth of dilute nitric acid followed by a few drops of aqueous silver nitrate. |
| | Record your observations. |
| | [1] |
| 4. | |
| (K) | Identify solid D . |
| | ro1 |
| | [2] |
| | [Total: 16] |

3 The label on a bottle of orange drink states 'contains no artificial colours'.

A scientist thinks that the orange colour in the drink is a mixture of two artificial colours:

- Sunset Yellow E110
- Allura Red E129.

Plan an experiment to show that the orange colour in the drink does **not** contain these two artificial colours.

Your plan should describe the use of common laboratory apparatus and samples of E110, E129 and the orange colouring from the drink.

You may draw a diagram to help answer the question.

| [6] |
|-----|

Notes for use in qualitative analysis

Tests for anions

| anion | test | test result |
|---|--|---|
| carbonate, CO ₃ ²⁻ | add dilute acid, then test for carbon dioxide gas | effervescence, carbon dioxide produced |
| chloride, C <i>l</i> ⁻ [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | white ppt. |
| bromide, Br ⁻ [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | cream ppt. |
| iodide, I ⁻ [in solution] | acidify with dilute nitric acid, then add aqueous silver nitrate | yellow ppt. |
| nitrate, NO ₃ ⁻ [in solution] | add aqueous sodium hydroxide, then aluminium foil; warm carefully | ammonia produced |
| sulfate, SO ₄ ²⁻ [in solution] | acidify with dilute nitric acid, then add aqueous barium nitrate | white ppt. |
| sulfite, SO ₃ ²⁻ | add a small volume of acidified aqueous potassium manganate(VII) | the acidified aqueous potassium manganate(VII) changes colour from purple to colourless |

Tests for aqueous cations

| cation | effect of aqueous sodium hydroxide | effect of aqueous ammonia |
|--|--|--|
| aluminium, Al ³⁺ | white ppt., soluble in excess, giving a colourless solution | white ppt., insoluble in excess |
| ammonium, NH ₄ ⁺ | ammonia produced on warming | - |
| calcium, Ca ²⁺ | white ppt., insoluble in excess | no ppt. or very slight white ppt. |
| chromium(III), Cr ³⁺ | green ppt., soluble in excess | green ppt., insoluble in excess |
| copper(II), Cu ²⁺ | light blue ppt., insoluble in excess | light blue ppt., soluble in excess, giving a dark blue solution |
| iron(II), Fe ²⁺ | green ppt., insoluble in excess, ppt. turns brown near surface on standing | green ppt., insoluble in excess, ppt. turns brown near surface on standing |
| iron(III), Fe ³⁺ | red-brown ppt., insoluble in excess | red-brown ppt., insoluble in excess |
| zinc, Zn ²⁺ | white ppt., soluble in excess, giving a colourless solution | white ppt., soluble in excess, giving a colourless solution |

Tests for gases

| gas | test and test result |
|---------------------------------|--|
| ammonia, NH ₃ | turns damp red litmus paper blue |
| carbon dioxide, CO ₂ | turns limewater milky |
| chlorine, Cl ₂ | bleaches damp litmus paper |
| hydrogen, H ₂ | 'pops' with a lighted splint |
| oxygen, O ₂ | relights a glowing splint |
| sulfur dioxide, SO ₂ | turns acidified aqueous potassium manganate(VII) from purple to colourless |

Flame tests for metal ions

| metal ion | flame colour |
|------------------------------|--------------|
| lithium, Li ⁺ | red |
| sodium, Na ⁺ | yellow |
| potassium, K ⁺ | lilac |
| calcium, Ca ²⁺ | orange-red |
| barium, Ba ²⁺ | light green |
| copper(II), Cu ²⁺ | blue-green |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.