CAMBRIDGE
International Education

Cambridge International AS & A Level

FURTHER MATHEMATICS **9231/04**

Paper 4 Further Probability & Statistics **For examination from 2027**

MARK SCHEME

Maximum Mark: 50

Specimen

This document has **10** pages.

© Cambridge University Press & Assessment 2026

General Marking Principles

All examiners must apply these marking principles when marking candidate responses.

1	Examiners must award marks for each response in line with: • the specific **content** defined in the mark scheme • the specific **skills** defined in the mark scheme or in the level descriptions • the **standard of response** required as exemplified by the standardisation scripts.
2	Examiners must award **whole marks** (not half marks, or other fractions).
3	Examiners must award marks **positively**. This means that: • marks are not deducted for errors or omissions • marks are awarded for correct/valid answers as defined in the mark scheme and also for **other valid answers** which go beyond the scope of the syllabus and mark scheme referring to your team leader as appropriate • the quality of spelling, punctuation and grammar are judged **only** when the mark scheme indicates that these features are **specifically assessed** in the question. However, the meaning of all responses must be unambiguous.
4	Examiners must **consistently** apply the requirements of the mark scheme and any rules for what to do when candidates have not followed instructions correctly.
5	Examiners must use the **full range of marks** defined in the mark scheme, unless limited by the quality of the candidate responses seen.
6	Examiners must award marks according to the mark scheme and **must not** award marks with grade thresholds or grade descriptions in mind.

© Cambridge University Press & Assessment 2026

Mathematics-Specific Marking Principles

1 Unless a particular method has been specified in the question, full marks may be awarded for any correct method. However, if a calculation is required then no marks will be awarded for a scale drawing.

2 Unless specified in the question, non-integer answers may be given as fractions, decimals or in standard form. Ignore superfluous zeros, provided that the degree of accuracy is not affected.

3 Allow alternative conventions for notation if used consistently throughout the paper, e.g. commas being used as decimal points.

4 Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored (isw).

5 Where a candidate has misread a number or sign in the question and used that value consistently throughout, provided that number does not alter the difficulty or the method required, award all marks earned and deduct just 1 A or B mark for the misread.

6 Recovery within working is allowed, e.g. a notation error in the working where the following line of working makes the candidate's intent clear.

© Cambridge University Press & Assessment 2026

Mark Scheme Notes

Marks are of the following three types.

M	Method mark, given for a valid method applied to the problem. Method marks can still be given even if there are numerical errors, algebraic slips or errors in units. However the method must be applied to the specific problem, e.g. by substituting the relevant quantities into a formula. Correct use of a formula without the formula being quoted earns the M mark and in some cases an M mark can be implied from a correct answer.
A	Accuracy mark, given for an accurate answer or accurate intermediate step following a correct method. Accuracy marks cannot be given unless the relevant method mark has also been given.
B	Mark for a correct statement or step.
DM or DB	M marks and B marks are generally independent of each other. The notation DM or DB means a particular M or B mark is dependent on an earlier M or B mark (indicated by *). When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.

- A or B marks are given for correct work only (not for results obtained from incorrect working) unless follow through is allowed (see abbreviation FT below).
- Wrong or missing units in an answer should not result in loss of marks unless the guidance indicates otherwise.
- For a numerical answer, allow the A or B mark if the answer is correct to 3 significant figures (sf) or would be correct to 3 sf if rounded (1 decimal place (dp) in the case of an angle in degrees). As stated above, an A or B mark is not given if a correct numerical answer is obtained from incorrect working.
- Common alternative solutions are shown in the Answer column as: '**EITHER** Solution 1 **OR** Solution 2 **OR** Solution 3 …'. Round brackets appear in the Partial Marks column around the marks for each alternative solution.
- The total number of marks available for each question is shown at the bottom of the Marks column in bold type.
- Square brackets [ ] around text show extra information not needed for the mark to be awarded.

The following abbreviations may be used in a mark scheme.

AG	Answer given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid).
CAO	Correct answer only (emphasising that no 'follow through' from an error is allowed).
CWO	Correct working only
FT	Follow through after error (see Mark Scheme Notes for further details).
ISW	Ignore subsequent working
OE	Or equivalent form
SC	Special case
SOI	Seen or implied

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Partial Marks	Guidance
1(a)	When the population cannot be assumed to be normally distributed.	1	**B1**	OE
		1		
1(b)	H_0: population median is 6.00, H_1: population median is greater than 6.00	1	**B1**	Both hypotheses stated.
	Calculate deviations and resulting signed ranks.	1	**M1**	
	<table><tr><td>**Devs**</td><td>–0.38</td><td>–0.27</td><td>0.55</td><td>0.81</td><td>0.10</td><td>–0.25</td><td>–0.13</td><td>0.47</td><td>–0.14</td><td>0.26</td><td>0.99</td><td>–0.09</td></tr><tr><td>**Rank**</td><td>–8</td><td>–7</td><td>10</td><td>11</td><td>2</td><td>–5</td><td>–3</td><td>9</td><td>–4</td><td>6</td><td>12</td><td>–1</td></tr></table>	1	**A1**	
	Test statistic $T = 8 + 7 + 5 + 3 + 4 + 1 = 28$	1	**A1**	
	Since $28 > 17$, accept H_0.	1	**M1**	Allow equivalent forms, for example do not reject H_0 or not significant.
	Insufficient evidence to suggest that the average pollution level in the river is greater than 6.00.	1	**A1**	Conclusion to be stated in context, not just ‘not significant’; follow through *their* value of T.
		6		

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Partial Marks	Guidance
2	H_0: Geo(0.6) is a satisfactory model for the data. H_1: Geo(0.6) is not a satisfactory model for the data.	1	**B1**	
	Use Geo(0.6) to calculate expected values 120, 48, 19.2, 7.68, 3.072, 1.2288, 0.8192	1	**M1**	Need at least 4 values correct
	Last 3 cells combined	1	**M1**	Observed frequency = 6, expected frequency = 5.12
	$\frac{(126-120)^2}{120}+\frac{(43-48)^2}{48}+\frac{(22-19.2)^2}{19.2}+\frac{(3-7.68)^2}{7.68}+\frac{(6-5.12)^2}{5.12}$	1	**M1**	0.3 + 0.5208 + 0.4083 + 2.8519 + 0.15125
	$\chi^2 = 4.23$	1	**A1**	Correct value to (at least) 3 significant figures.
	Since $4.23 < 9.49$, accept H_0.	1	**M1**	Allow equivalent forms, for example do not reject H_0 or not significant.
	Insufficient evidence to suggest that Geo(0.6) is not a satisfactory model (for the data).	1	**A1**	Correct conclusion from correct values; conclusion must refer to the distribution explicitly.
		7		

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Partial Marks	Guidance
3	Assume that population differences are normally distributed.	1	**B1**	
	$H_0 : \mu_B - \mu_A = 0$, $H_1 : \mu_B - \mu_A > 0$	1	**B1**	Both hypotheses required, in terms of population means or alternatively in terms of population mean difference.
	Differences, d: 8 3 –4 2 –11 3 17 10 6 0	1	**M1**	
	$\overline{d} = 3.4$ and $s^2 = \frac{1}{9}\left(648 - \frac{34^2}{10}\right) = 59.16$	1	**M1**	
	Calculate test statistic $t = \frac{\overline{d}}{s/\sqrt{10}}$	1	**M1**	
	$t = 1.398$	1	**A1**	Correct value to at least 3 significant figures.
	Since $1.398 > 1.383$, reject H_0.	1	**M1**	
	Sufficient evidence to suggest that the population mean number of hours of absence has decreased following the introduction of flexi-time.	1	**A1**	
		8		

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Partial Marks	Guidance
4	Sample variances are $\frac{4341.6}{59}$ and $\frac{3732.0}{49}$	1	**M1**	Either attempted.
	$s_x^2 = 73.59$ and $s_y^2 = 76.16$	1	**A1**	Both correct to at least 2 decimal places.
	Variance for difference of means: $s^2 = \frac{s_x^2}{60} + \frac{s_y^2}{50}$	1	**M1**	
	$s^2 = 2.750$ or $s = 1.658$	1	**A1**	Correct value to at least 4 significant figures.
	Evaluate $(\overline{x} - \overline{y}) \pm zs$	1	**M1**	Any z-value, and using *their* value of s.
	Use of correct $z = 1.96$	1	**A1**	
	[1.55, 8.05]	1	**A1**	Accept e.g. (1.55, 8.05) or $1.55 < \mu_A - \mu_B < 8.05$.
		7		

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Partial Marks	Guidance
5(a)	$P(X > 1) = 1 - \frac{1}{2} \times 1 \times \frac{6}{5} = \frac{2}{5}$ **OR** via $\int_1^{\infty} \frac{6}{5}x^{-4}\,dx$	1	**B1**	
		1		
5(b)	Median satisfies $\frac{1}{2} \times m \times \frac{6}{5}m = \frac{1}{2}$ **OR** equivalent use of integration	1	**M1**	
	$m = \sqrt{\frac{5}{6}}$	1	**A1**	Correct value; accept 0.913.
		2		
5(c)	$E(X^2) = \int_0^1 \frac{6}{5}x^3\,dx + \int_1^{\infty} \frac{6}{5}x^{-2}\,dx$	1	**M1**	Attempt evaluation of the sum of these two integrals.
	$E(X^2) = 1.5$	1	**A1**	Correct value seen or implied.
	$\text{Var}(X) = 1.5 - 1^2 = 0.5$	1	**A1FT**	Follow through *their* 1.5 providing variance is positive.
		3		
5(d)	$E(\sqrt{X}) = \int_0^1 \frac{6}{5}x^{\frac{3}{2}}\,dx + \int_1^{\infty} \frac{6}{5}x^{-\frac{7}{2}}\,dx$	1	**M1**	Attempt evaluation of the sum of these two integrals.
	$\frac{24}{25}$ or 0.96	1	**A1**	
		2		

© Cambridge University Press & Assessment 2026

Question	Answer	Marks	Partial Marks	Guidance
6(a)	Probabilities $\frac{1}{8}, \frac{3}{8}, \frac{3}{8}, \frac{1}{8}$ for 0, 1, 2, 3 reds	1	**B1**	
	$G_X(t) = \frac{1}{8} + \frac{3}{8}t + \frac{3}{8}t^2 + \frac{1}{8}t^3$	1	**B1FT**	Follow through *their* probabilities so long as $\Sigma p = 1$.
		2		
6(b)	Correct value $\frac{9}{20}$ or 0.45 for $P(Y = 1)$ or $P(Y = 2)$	1	**B1**	
	Probabilities $\frac{1}{20}, \frac{9}{20}, \frac{9}{20}, \frac{1}{20}$ for 0, 1, 2, 3 reds	1	**B1**	
	$G_Y(t) = \frac{1}{20} + \frac{9}{20}t + \frac{9}{20}t^2 + \frac{1}{20}t^3$	1	**B1FT**	Follow through *their* probabilities so long as $\Sigma p = 1$.
		3		
6(c)(i)	$G_Z(t) = G_X(t) \times G_Y(t)$	1	**B1**	Stated or implied.
	Complete expansion of the product of *their* cubics.	1	**M1**	
	$G_Z(t) = \frac{1}{160}(1 + 12t + 39t^2 + 56t^3 + 39t^4 + 12t^5 + t^6)$	1	**A1**	Or equivalent, e.g. with explicit fractional coefficients.
		3		
6(c)(ii)	Attempt to differentiate $G_Z(t)$ and evaluate $G'_Z(1)$.	1	**M1**	
	$E(Z) = 3$	1	**A1**	Correctly obtained from $G'_Z(1)$.
	Attempt to find second derivative $G''_Z(t)$.	1	**M1**	
	Use of $G''_Z(1) + G'_Z(1) - \left(G'_Z(1)\right)^2$.	1	**M1**	
	$\text{Var}(Z) = 1.2$	1	**A1**	Correct value correctly obtained.
		5		

© Cambridge University Press & Assessment 2026